

400 mm	
HIGHLIGHTS OF PRESCRIBING INFORMATION	
These highlights do not include all the information needed to use NICARDIPINE HYDROCHLORIDE IN SODIUM CHLORIDE INJECTION safely and effectively. See full prescribing information for NICARDIPINE HYDROCHLORIDE IN SODIUM CHLORIDE INJECTION. NICARDIPINE HYDROCHLORIDE IN SODIUM CHLORIDE INJECTION, for intravenous use	
Initial U.S. Approval: 1988	
Rx Only	
-----RECENT MAJOR CHANGES-----	
Dosage Forms and Strengths (3)	08/2024
-----INDICATIONS AND USAGE-----	
Nicardipine hydrochloride in 0.9% sodium chloride injection is a calcium channel blocker indicated for the short-term treatment of hypertension when oral therapy is not feasible. (1)	
-----DOSAGE AND ADMINISTRATION-----	
• Individualize dosage based upon the severity of hypertension and response of the patient during dosing (2, 1). • When substituting for oral nicardipine therapy, use the intravenous infusion rate as follows (2, 3):	
Oral Nicardipine Dose	Equivalent Intravenous Infusion Rate
20 mg every 8 hours	0.5 mg/hr
30 mg every 8 hours	1.2 mg/hr
40 mg every 8 hours	2.2 mg/hr

	FULL PRESCRIBING INFORMATION: CONTENTS*
	1 INDICATIONS AND USAGE
	1.1 Hypertension
	2 DOSAGE AND ADMINISTRATION
	2.1 General Information
	2.2 Inspection and Preparation
	2.3 Dosage as a Substitute for Oral Nifedipine Therapy
	2.4 Dosage for Initiation of Therapy in a Drug-Free Patient
	2.5 Conditions Requiring Intensive Adjustment
	2.6 Transition to Oral Antihypertensive Agents
	3 DOSAGE FORMS AND STRENGTHS
	4 CONTRAINDICATIONS
	4.1 Advanced Aortic Stenosis
	5 WARNINGS AND PRECAUTIONS
	5.1 Excessive Pharmacologic Effects
	5.2 Rapid Decreases in Blood Pressure
	5.3 Use in Patients with Angina
	5.4 Use in Patients with Congestive Heart Failure
	5.5 Use in Patients with Impaired Hepatic Function
	5.6 Use in Patients with Impaired Renal Function
	5.7 Intravenous Infusion Site
	5.8 Beta-Blocker Withdrawal
	5.9 Use in Patients with Pheochromocytoma
	6 ADVERSE REACTIONS
	6.1 Adverse Reactions Observed in Clinical Trials
	7 DRUG INTERACTIONS
	7.1 Antihypertensive Agents

FULL PRESCRIPTION INFORMATION

1. INDICATIONS AND USAGE

1.1 Hypertension

Nicardipine hydrochloride in 0.9% sodium chloride injection is indicated for the short-term treatment of hypertension when oral therapy is not feasible or desirable. For optimal control of blood pressure, transfer patients to oral medication as soon as their clinical condition permits (see Dosage and Administration 2.6a).

2. DOSAGE AND ADMINISTRATION

2.1 General Information

Individual dosing based on the severity of hypertension and the response of the patient during dosing. Monitor blood pressure and heart rate both during and after infusion to avoid tachycardia or too rapid or excessive reduction in either systolic or diastolic blood pressure.

Administer nicardipine hydrochloride in 0.9% sodium chloride injection by slow continuous infusion by a central line or through a large peripheral vein. Change the infusion site every 12 hours if administered via peripheral vein [see Intravenous Infusion Site (5.7)].

2.2 Inspection and Preparation

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use the solution if particulate matter, precipitate, or crystallization is present, or if the container appears damaged.

Single Dose Contains

Dilution is not required for nifedipine hydrochloride in 0.9% sodium chloride injection.

Check the container for minute leaks just prior to use; ensure that the seal is intact. If leaks are found, discard solution as sterility may be impaired.

Do not use nifedipine hydrochloride in 0.9% sodium chloride injection with any product in the same intravenous line or premixed container. Do not add supplementary medication to the bag. Protect from light until ready to use.

Do not use plastic containers in series connections. Use could result in air embolism due to residual air being drawn from the primary container before the administration of the fluid from the secondary container is complete.

Discard Unused Portion

Preparation for administration

1. Suspend container from eyelinet support.
2. Remove protection from outlet port of bottom of container.
3. Attach administration set. Refer to complete directions accompanying set.

2.3 Dose as a Substitute for Oral Nifedipine

The intravenous infusion rate required to produce an average plasma concentration equivalent to a given oral dose at steady state is shown in the following table:

Oral Nifedipine Dose	Equivalent Intravenous Infusion Rate
20 mg every 8 hours	0.5 mg/hr
30 mg every 8 hours	1.2 mg/hr
40 mg every 8 hours	2.2 mg/hr

2.4 Dose for Initiation of Therapy in a Drug-Free Patient

The time course of blood pressure decrease is dependent on the initial rate of infusion. The frequency of dose adjustment. Nifedipine hydrochloride in 0.9% sodium chloride injection is administered by slow continuous infusion at a concentration of 0.1 mg/mL. With constant infusion, blood pressure begins to fall within 10 minutes. It reaches about 50% of its ultimate decrease in about 45 minutes.

When treating acute hypertensive episodes in patients with chronic hypertension, discontinuation of infusion is followed by a 50% offset of action in 30 minutes plus 5 minutes at plasma levels of drug and slightly decreasing antihypertensive effect for many hours.

ARTWORK INFORMATION	
Item Code: 22201534	
Component : Leaflet	
Dimension: Length - 920, Height - 450 mm	
Customer Name: Caplin Steriles	
Specification Substrate: 40 GSM Bible Paper	
Language: English	Country: US
Change History: New Artwork	Colours: 1 BLACK
	Die Line :

- In a drug-free patient, initiate therapy at 5 mg/hr. Increase the infusion rate by 2.5 mg/hr to a maximum of 15 mg/hr until desired blood pressure reduction is achieved. For a gradual blood pressure reduction the rate can be increased every 15 minutes for a rapid reduction, every 5 minutes (2,4). - If hypotension or tachycardia ensues, discontinue the infusion. After stabilized, patient can be restarted at low doses such as 3 mg/hr to 5 mg/hr (2,5).	
-----DOSAGE FORMS AND STRENGTHS-----	
20 mg nifedipine hydrochloride in 200 mL 0.9% sodium chloride injection (0.1 mg/mL) in a single dose container (3) 40 mg nifedipine hydrochloride in 200 mL 0.9% sodium chloride injection (0.2 mg/mL) in a single container (3)	
-----CONTRAINDICATIONS-----	
Do not use in patients with advanced aortic stenosis (4,1).	
-----WARNINGS AND PRECAUTIONS-----	
- To reduce the possibility of venous thrombosis, phlebitis, and vascular impairment, do not use small veins, such as those on the dorsum of the hand or wrist. Avoid intraarterial administration or extravasation (5,7). - To minimize the risk of peripheral venous irritation, change the site of infusion of nifedipine every 12 hours (5,7). - Nifedipine is not a beta-blocker and therefore gives no protection against the dangers of abrupt beta-blocker withdrawal. Withdraw beta-blockers gradually (5,8). - Closely monitor response in patients with angina (5,3), congestive heart failure (5,4), impaired hepatic function (5,5), portal hypertension (5,6), and renal impairment (5,6) and pheochromocytoma (5,9).	
-----ADVERSE REACTIONS-----	
Most common adverse reactions are headache (13%), hypotension (5%), tachycardia (4%) and nausea/vomiting (4%). (6)	
-----DRUG INTERACTIONS-----	
- Cimetidine increases nifedipine plasma levels (7,3). - Nifedipine may increase cyclosporine and tacrolimus plasma levels. Frequent monitoring of trough blood levels of cyclosporine and tacrolimus is recommended when co-administering nifedipine. (7,5,7,6).	
-----USE IN SPECIFIC POPULATIONS-----	
- Pregnancy: Based on animal data, may cause fetal harm (6,1). - Nursing Mothers: It is recommended that women who wish to breastfeed should not be given this drug (8,3). - Safety and efficacy in patients under the age of 18 have not been established (8,4).	
Revised: 11/2023	
7.2	Beta-Blockers
7.3	Cimetidine
7.4	Digoxin
7.5	Cyclosporine
7.6	Tacrolimus
7.7	In Vitro Interaction
8	USE IN SPECIFIC POPULATIONS
8.1	Pregnancy
8.3	Nursing Mothers
8.4	Pediatric Use
8.5	Geriatric Use
10	OVERDOSAGE
11	DESCRIPTION
12	CLINICAL PHARMACOLOGY
12.1	Mechanism of Action
12.2	Pharmacodynamics
12.3	Pharmacokinetics
13	NONCLINICAL TOXICOLOGY
13.1	Carcinogenesis, Mutagenesis, Impairment of Fertility
13.3	Reproductive and Developmental Toxicology
14	CLINICAL STUDIES
16	HOW SUPPLIED/STORAGE AND HANDLING
16.1	How Supplied
16.2	Storage and Handling

*Sections or subsections omitted from the full prescribing information are not listed.

Indications

For a gradual reduction in blood pressure, initiate therapy at a rate of 5 mg/hr. If desired blood pressure reduction is not achieved at this dose, increase the infusion rate by 2.5 mg/hr every 15 minutes up to a maximum of 15 mg/hr, until desired blood pressure reduction is achieved. For more rapid blood pressure reduction, titrate every 5 minutes.

Maintenance

Adjust the rate of infusion as needed to maintain desired response.

2.5 Conditions Requiring Infusion Adjustment

Hypotension or Tachycardia:

In case of hypotension or tachycardia, discontinue infusion. When blood pressure and heart rate stabilize, restart infusion at low doses such as 30 mL/hr to 50 mL/hr (3 mg/hr to 5 mg/hr) and titrate to maintain desired blood pressure.

Infusion Site Changes:

Change infusion site every 12 hours if administered via peripheral vein.

Impaired Cardiac, Hepatic, or Renal Function:

Monitor closely when titrating nicardipine hydrochloride in 0.9% sodium chloride injection in patients with congestive heart failure or impaired hepatic or renal function [see Warnings and Precautions (5.4, 5.5 and 5.6)].

2.6 Transfer to Oral Antihypertensive Agents

Treatment includes transfer to an oral antihypertensive agent other than nicardipine capsules, initiate oral therapy upon discontinuation of nicardipine hydrochloride in 0.9% sodium chloride injection.

When switching to a three times a day regimen of nicardipine capsules, administer the first dose 1 hour prior to discontinuation of the infusion.

3 DOSAGE FORMS AND STRENGTHS

Nicardipine hydrochloride in 0.9% sodium chloride injection is a clear, colorless to yellow solution and is available in the following presentations:

- 20 mg nicardipine hydrochloride in 200 mL 0.9% sodium chloride injection (0.1 mg/mL) in a single dose container
- 40 mg nicardipine hydrochloride in 200 mL 0.9% sodium chloride injection (0.2 mg/mL) in a single container

4 CONTRAINDICATIONS

4.1 Advanced Aortic Stenosis

Do not use nicardipine in patients with advanced aortic stenosis because of the afterload reduction effect of nicardipine. Reduction of diastolic pressure in these patients may worsen rather than improve myocardial oxygen balance.

5 WARNINGS AND PRECAUTIONS

5.1 Excessive Pharmacologic Effects

In administering nicardipine, close monitoring of blood pressure and heart rate is required. Nicardipine may occasionally produce symptomatic hypotension or tachycardia. Avoid systemic hypotension when administering the drug to patients who have sustained an acute cerebral infarction or hemorrhage.

5.2 Rapid Decreases in Blood Pressure

No clinical events have been reported suggestive of a too rapid decrease in blood pressure with nicardipine. However, as with any antihypertensive agent, blood pressure lowering should be accomplished over as long a time as is compatible with the patient's clinical status.

5.3 Use in Patients with Angina

Increases in frequency, duration, or severity of angina have been seen in chronic oral therapy with nicardipine capsules. Induction or exacerbation of angina has been seen in less than 1% of coronary artery disease patients treated with nicardipine capsules. The mechanism of this effect has not been established.

5.4 Use in Patients with Congestive Heart Failure

Nicardipine reduced afterload without impairing myocardial contractility in preliminary hemodynamic studies of CHF patients. However, in vitro and in some patients, a negative inotropic effect has been observed. Therefore, monitor vital signs carefully when using nicardipine, particularly in combination with a beta-blocker, in patients with CHF or significant left ventricular dysfunction.

5.5 Use in Patients with Impaired Hepatic Function

Since nicardipine is metabolized in the liver, consider lower dosages and closely monitor response. Nicardipine administered intravenously increased hepatic venous pressure gradient by 4 mmHg in cirrhotic patients at high doses (5 mg/20 min) in one study. Use caution in patients with portal hypertension.

5.6 Use in Patients with Impaired Renal Function

When nicardipine was given to mild-to-moderate hypertensive patients with moderate renal impairment, a significantly lower systemic clearance and higher AUC was observed. These results are consistent with those seen after oral administration of other antihypertensive drugs. Careful dose titration is advised when treating patients with more than mild renal impairment.

5.7 Intravenous Infusion Site

To reduce the possibility of venous thrombosis, phlebitis, local irritation, swelling, extravasation, and the rare occurrence of vascular impairment, administer drug through large peripheral veins or central veins rather than arteries or small peripheral veins, such as those on the dorsum of the hand or wrist. Minimize the risk of peripheral venous irritation, consider changing the site of the drug infusion every 12 hours.

5.8 Beta-Blocker Withdrawal

Nicardipine is not a beta-blocker and therefore gives no protection against the dangers of abrupt beta-blocker withdrawal. Withdraw beta-blockers gradually.

5.9 Use in Patients with Pheochromocytoma

Only limited clinical experience exists in use of nicardipine for patients with hypertension from pheochromocytoma.

920 mm

6 ADVERSE REACTIONS

6.1 Adverse Reactions Observed in Clinical Trials

Because clinical trials are conducted under widely varying conditions, adverse reaction rates observed in the clinical trials of a drug cannot be directly compared to rates in the clinical trials of another drug and may not reflect the rates observed in practice.

Two hundred forty-four patients participated in a multicenter, double-blind, placebo-controlled trials of nicanipride. Adverse experiences were generally not serious and most were expected consequences of vasodilation. Adverse reactions occasionally required dose adjustment. Therapy was discontinued in approximately 12% of patients, mainly due to hypotension, headache, and tachycardia. Adverse reactions that occurred more often on nicanipride than on placebo by at least 2% were headache (12%) and nausea/vomiting (4%).

The following adverse reactions have been reported in clinical trials or in the literature during the use of intravenously administered nicanipride.

Body as a Whole: fever, neck pain
Cardiovascular: angina pectoris, atrioventricular block, ST segment depression, inverted T wave, deep-vein thrombophlebitis
Digestive: dyspepsia
Hemic and Lymphatic: thrombocytopenia
Metabolic and Nutritional: hypophosphatemia, peripheral edema
Nervous: confusion, hypertension
Respiratory: respiratory disorder
Special Senses: conjunctivitis, ear disorder, tinnitus
Urogenital: urinary frequency

Sinus node dysfunction and myocardial infarction, which may be due to disease progression, have been seen in patients on chronic therapy with orally administered nicanipride.

7 DRUG INTERACTIONS

7.1 Antihypertensive Agents

Since nicanipride hydrochloride injection may be administered simultaneously with other medications, including other antihypertensive agents, careful monitoring of these patients is necessary to detect and to treat promptly any undesired effects from concomitant administration.

7.2 Beta-Blockers

In most patients, nicanipride hydrochloride injection can safely be used concomitantly with beta-blockers. However, monitor response carefully when combining nicanipride hydrochloride injection with a beta-blocker in the treatment of congestive heart failure patients (see Warnings and Precautions (5.4)).

7.3 Cimetidine

Cimetidine has been shown to increase nicanipride plasma concentrations with oral nicanipride administration. Carefully monitor patients receiving the two drugs concomitantly. Data with other histamine-2 antagonists are not available.

7.4 Digoxin

Studies have shown that oral nicanipride usually does not alter digoxin plasma concentrations.

7.5 Cyclosporine

Concomitant administration of oral or intravenous nicanipride and cyclosporine results in elevated plasma cyclosporine levels through nicanipride inhibition of hepatic microsomal enzymes, including CYP3A4. Monitor closely plasma concentrations of cyclosporine during nicanipride hydrochloride injection administration, and adjust the dose of cyclosporine accordingly.

7.6 Tacrolimus

Concomitant administration of intravenous nicanipride and tacrolimus may result in elevated plasma tacrolimus levels through nicanipride inhibition of hepatic microsomal enzymes, including CYP3A4. Closely monitor plasma concentrations of tacrolimus during nicanipride administration, and adjust the dose of tacrolimus accordingly.

7.7 In Vitro Interaction

The plasma protein binding of nicanipride was not altered when therapeutic concentrations of furosemide, propranolol, dipyridamide, warfarin, quinine, or naproxen were added to human plasma *in vitro*.

8 USE IN SPECIFIC POPULATIONS

8.1 Pregnancy

Pregnancy Category C.

There are no adequate and well-controlled studies of nicanipride use in pregnant women. There are limited human data in pregnant women with pre-eclampsia and preterm labor. In animal reproduction and developmental toxicity studies, evidence of fetal harm was observed. Therefore use nicanipride during pregnancy only if the potential benefit justifies the potential risk to the fetus.

In reproduction studies conducted in rats and rabbits, increased embryofetality occurred when nicanipride was administered intravenously at doses equivalent to human intravenous doses of 1.6 (rats) and 0.32 mg/kg/day (rabbits). Increased embryofetality was also observed when nicanipride was administered orally to pregnant rabbits at a dose equivalent to a human oral dose of about 48 mg/kg/day in dose 24 times the maximum recommended human oral dose and once daily for 14 days. In rabbits, maternal body weight was suppressed. At a lower oral dose, equivalent to a human dose of about 32 mg/kg/day (16 times the maximum recommended human oral dose), in a different strain of rabbit, there were no adverse effects on the fetus, though there was increased maternal mortality. There was no evidence of embryofetality or teratogenicity when pregnant rats were administered nicanipride orally at a dose equivalent to a human oral dose of about 16 mg/kg/day (8 times the MRHD); however, oocyte, reduced birth weight, reduced neonatal survival and reduced neonatal weight gain were reported (see Nonclinical Toxicology (13.3)).

8.3 Nursing Mothers

Nicanipride is minimally excreted into human milk. Among 18 infants exposed to nicanipride through breast milk in the postpartum period, calculated daily infant dose was less than 0.3 mcg and there were no adverse events observed. It is not known whether women who wish to breastfeed should not be given this drug.

In a study of 11 women who received oral nicanipride 4 days to 14 days postpartum, 4 women received immediate-release nicanipride 40 to 80 mg daily, 6 women received sustained-release nicanipride 100 mg to 150 mg daily, and one woman received

the analysis of seven women who received intravenous nicardipine for an average of 1.9 days in the immediate postpartum period as therapy for pre-eclampsia, 34 milk samples were obtained at unspecified times and nicardipine was undetectable (less than 5 mcg/L) in 82% of the samples. Four women who received 1 to 6.5 mcg/hour of nicardipine had 6 milk samples with detectable nicardipine levels (range 5.1 to 18.5 mcg/L). The highest concentration of 18.5 mcg/L was found in a woman who received 5.5 mcg/hour of nicardipine. The estimated maximum dose in a breastfed infant was less than 0.3 mcg daily or 0.015% to 0.004% of the therapeutic dose in a 1 kg infant.

8.4 Pediatric Use
Safety and efficacy in patients under the age of 18 have not been established.

8.5 Geriatric Use
The steady-state pharmacokinetics of nicardipine are similar in elderly hypertensive patients (greater than 65 years) and young healthy adults. Clinical studies of nicardipine did not include sufficient numbers of subjects aged 65 and over to determine whether they respond differently from younger subjects. Other reported clinical experience has not identified differences in responses between the elderly and younger patients. In general, dose selection for an elderly patient should be cautious, usually starting at the low end of the dosing range, reflecting the greater frequency of decreased hepatic, renal or cardiac function, and concomitant disease or other drug therapy.

9 OVERDOSEAGE
Several overdosages with orally administered nicardipine have been reported. One adult patient allegedly ingested 600 mg of nicardipine immediate release capsules, and another patient, 2160 mg of the sustained release formulation of nicardipine. Symptoms included marked hypotension, bradycardia, palpitations, flushing, drowsiness, confusion and slurred speech. All symptoms resolved without sequelae. An overdose occurred in a one-year-old child who ingested half of the powder in a 30 mg nicardipine standard capsule. The child remained asymptomatic. Based on results obtained in laboratory animals, lethal overdose may cause systemic hypotension, bradycardia (following initial tachycardia) and progressive atrioventricular conduction block. Reversible hepatic function abnormalities and sporadic focal hepatic necrosis were noted in some animal species receiving very large doses of nicardipine.

For treatment of overdose, standard measures including monitoring of cardiac and respiratory functions should be implemented. The patient should be positioned to avoid cerebral anoxia. Reverse blood pressure determinations are essential. Vasopressors are clinically indicated to prevent hypotension. Intravenous calcium gluconate may help reverse the effects of calcium entry blockade.

11 DESCRIPTION
Nicardipine hydrochloride is a calcium ion influx inhibitor (slow channel blocker or calcium channel blocker). Nicardipine hydrochloride is a dihydropyridine derivative with IUPAC (International Union of Pure and Applied Chemistry) chemical name: (±)-2-benzyl-2-methyl amino ethyl methyl 1,4-dihydro-2,6-dimethyl-4-[(m-nitrophenyl)-3,5-pyridinedicarboxylate] monohydrochloride and has the following structure:

CN(C)CCc1ccc(cc1)C2=C(C(=O)OC)C(=CN2C3=CC=CC=C3)C4=CC(=CC=C4)C(=O)OCC5C(C)=CC(=CC=C5)N

$\text{C}_{28}\text{H}_{28}\text{N}_2\text{O}_6 \cdot \text{HCl}$

Nicardipine hydrochloride is a yellow to pale yellow, odorless, crystalline powder with a melting-point range of 157-170 °C. It is soluble in methanol, sparingly soluble in ethanol, slightly soluble in acetone, chloroform and water. It has a molecular weight of 515.99.

Nicardipine hydrochloride in a 0.9% sodium chloride injection is available as a sterile, single-use, unit-of-use, iso-osmotic, clear, colorless to yellow solution for intravenous administration in a 200 mL single dose container. Each mL contains 0.1 mg of nicardipine hydrochloride. USP 9 mg sodium chloride. USP Hydrochloric acid (q.s.) may have been added to adjust pH to 3 to 5.

12 CLINICAL PHARMACOLOGY

12.1 Mechanism of Action

Nicardipine inhibits the transmembrane influx of calcium ions into cardiac muscle and smooth muscle without changing serum calcium concentrations. The contractile processes of cardiac muscle and vascular smooth muscle are dependent upon the movement of extracellular calcium ions into these cells through specific ion channels. The effects of nicardipine are more selective to vascular smooth muscle than cardiac muscle. In animal models, nicardipine produced relaxation of coronary vascular smooth muscle at drug levels which are in the negative inotropic effect.

12.2 Pharmacodynamics

Hemodynamics

Nicardipine produces significant decreases in systemic vascular resistance. In a study of intra-arterially administered nicardipine, the degree of vasodilation and the resultant decrease in blood pressure were more prominent in hypertensive patients than in normotensive volunteers. Administration of nicardipine to normotensive volunteers at dosages of 0.25 to 3 mg/hr for eight hours produced changes of less than 5 mmHg in systolic blood pressure and less than 3 mmHg in diastolic blood pressure.

An increase in heart rate is a normal response to vasodilation and decrease in blood pressure; in some patients these increases in heart rate may be pronounced. In placebo-controlled trials, the mean increases in heart rate were 7 ± 1 bpm in postoperative patients and 8 ± 1 bpm in patients with severe hypertension at the end of the maintenance period.

Hemodynamic studies following intravenous dosing in patients with coronary artery disease and normal or moderately abnormal left ventricular function have shown significant increases in ejection fraction and cardiac output with 1 mg/min infusion at a result decrease in left ventricular end-diastolic pressure (LVEDP). There is evidence that nicardipine increases blood flow. Coronary dilated infused by nicardipine improves peripheral and aortic metabolism in cases with chronic ischemia, resulting in reduced lactate production and augmented oxygen consumption. In patients with coronary artery disease, nicardipine, administered after beta-blockade, significantly improved systolic and diastolic left ventricular function. In congestive heart failure patients with impaired left ventricular function, nicardipine increased cardiac output both at rest and during exercise. Decreases in left ventricular end-diastolic pressure were also observed. However, in some patients with severe left ventricular dysfunction, it may have a negative inotropic effect and could lead to worsened failure.

"Coronary steal" has not been observed during treatment with nicardipine. Coronary steal is the detrimental redistribution of coronary blood flow in patients with coronary artery disease from underperfused areas toward better perfused areas.) Nicardipine has been shown to improve systolic shortening in both normal and hypokinetic segments of myocardial muscle. Radionucleid angiography has confirmed that wall motion remained improved during increased oxygen demand. (Occasional patients have developed increased angina upon receiving nicardipine capsules. Whether this represents coronary steal in these patients, or is the result of increased heart rate and decreased diastolic pressure, is not clear.)

In patients with coronary artery disease, nicardipine improves left ventricular diastolic distensibility during the early filling phase, probably due to a faster rate of myocardial relaxation in previously underperfed areas. There is little or no effect on normal myocardium, suggesting the improvement is mainly by indirect mechanisms such as afterload reduction and reduced ischemia. Nicardipine has no negative effect on myocardial relaxation at therapeutic doses. The clinical benefits of these properties have yet not been demonstrated.

Electrophysiology Effects

In general, no detrimental effects on the cardiac conduction system have been seen with nicardipine. During acute electrophysiologic studies, it prolonged heart rate and prolonged the corrected QT interval to a minor degree. It did not affect sinus node recovery or SA conduction times. The PA, AH, and HV intervals^a or the functional and effective refractory periods of the atrium were not increased. The relative and effective refractory periods of the His-Purkinje system were slightly shortened.^b A -conduction time was found with low to high right atrial; AH - conduction time from low right atrial to His bundle deflection; or AV nodal conduction time; H+V - conduction time through the His bundle and the bundle branch-Purkinje system.

Hepatic Function

Nicardipine hydrochloride is extensively metabolized by the liver. Plasma concentrations are influenced by changes in hepatic function. In a clinical study with nicardipine capsules in patients with severe liver disease, plasma concentrations were elevated and the half-life was prolonged [see Warnings and Precautions (5.5)]. Similar results were obtained in patients with hepatic disease when nicardipine hydrochloride was administered for 24 hours at 0.6 mg/hr.

Renal Function

When nicardipine was given to mild-to-moderate hypertensive patients with moderate degrees of renal impairment, significant reductions in glomerular filtration rate (GFR) and effective renal plasma flow (RPF) were observed. No significant differences in their flow rate were observed in these patients. A significantly lower systemic clearance and higher area under the curve (AUC) were observed.

When nicardipine capsules (20 or 30 mg TID) were given to hypertensive patients with impaired renal function, mean plasma concentrations, AUC, and C_{max} were approximately two-fold higher than in healthy controls. There was a transient increase in electrolyte excretion, including sodium [see Warnings and Precautions (5.6)]. Acute bolus administration of nicardipine hydrochloride injection (2.5 mg) in healthy volunteers decreased mean arterial pressure and renal vascular resistance; glomerular filtration rate (GFR), renal plasma flow (RPF), and the filtration fraction were unchanged. In healthy patients undergoing abdominal RFPI, nicardipine hydrochloride injection (10 mg over 20 minutes) increased GFR with no change in RPF when compared with placebo. In hypertensive type II diabetic patients with nephropathy, nicardipine hydrochloride (20 mg TID) did not change RPF and GFR, but reduced renal vascular resistance.

Pain Management

In well-controlled studies of patients with obstructive airway disease treated with nicardipine capsules, no evidence of increased bronchospasm was seen. In one of the studies, nicardipine capsules improved forced expiratory volume 1 second (FEV₁) and forced vital capacity (FVC) in combination with metoprolol. Adverse reactions reported in a limited number of patients with asthma, reactive airway disease, or obstructive airway disease were similar to reactions in other patients treated with nicardipine capsules.

12.3 Pharmacokinetics

Distribution

A rapid dose-related increase in nicardipine plasma concentrations is seen during the first two hours after the start of an infusion of nicardipine. Plasma concentrations increase at a much slower rate after the first few hours, and approach steady-state levels at 24 to 48 hours. The steady-state pharmacokinetics of nicardipine are similar in elderly hypertensive patients (greater than 65 years) and young healthy adults. On termination of the infusion, nicardipine concentrations decrease rapidly, with a half-life at least a 50% decrease during the first two hours post-infusion. The effects of nicardipine on plasma presence significantly correlate with plasma concentrations. Nicardipine is highly protein bound (greater than 95%) in human plasma over a wide concentration range.

After intravenous infusion, nicardipine plasma concentrations decline triexponentially, with a rapid early distribution phase ($t_{1/2}$ -half life of 3 minutes), an intermediate phase ($t_{1/2}$ -half life of 45 minutes), and a slow terminal phase ($t_{1/2}$ -half life of 14 hours). Data can only be detected after long-term infusions. Total plasma clearance (Cl) is 0.4 L/h/mg/kg, and the apparent volume of distribution (VD) using a non-compartment model is 8.3 L/kg. The pharmacokinetics of nicardipine is linear over the dosage ranges of 0.5 mg/kg to 40 mg/hr.

Metabolism and Excretion

Nicardipine has been shown to be rapidly and extensively metabolized by the hepatic cytochrome P450 enzymes, CYP2D6, 2D6, and 3A4. Nicardipine does not induce or inhibit its own metabolism; however, nicardipine has been shown to inhibit certain cytochrome P450 enzymes including CYP3A4, CYP2D6, CYP2C8, and CYP2C19. Inhibition of these enzymes may result in increased plasma levels of certain drugs, including cyclosporine and tacrolimus [see Drug Interactions (7.5, 7.6)]. The altered pharmacokinetics may necessitate dosage adjustment of the affected drug or discontinuation of treatment.

After coadministration of a radioactive intravenous dose of nicardipine with oral propanolol, 30% of the radioactivity was recovered in the urine and 43% in the feces within 96 hours. None of the dose was recovered as unchanged nicardipine.

13 UNLONCLINICAL TOXICOLOGY

13.1 Carcinogenesis, Mutagenesis, Impairment of Fertility

Rat treated with nicardipine in the diet at concentrations calculated to provide daily dosage levels of 5, 15, or 45 mg/kg/day for two years showed a dose-dependent increase in thyroid hyperplasia and neoplasia (follicular adenoma/carcinomas). One-month and three-month studies in the rat have suggested that these results are linked to a thyroid-independent reduction in plasma Thyroxine (T4) levels with a consequent increase in plasma levels of thyroid stimulating hormone (TSH). Chronic oral dosing of TSH to known potent cause hyperstimulation of the thyroid.

Rats on a low iodine deficient diet, nicardipine administration for one month was associated with thyroid hyperplasia that was prevented by T4 supplementation. Mice treated with nicardipine in the diet (at concentrations calculated to provide daily dosage levels up to 100 mg/kg/day for up to 18 months showed no evidence of neoplasia or any gross changes.

There was no evidence of thyrotoxic potential in dogs treated up to 25 mg nicardipine/kg/day for one year and no evidence of effects of nicardipine on thyroid function plasma T4 and TSH in man.

There was no occurrence of a mutagenic pathway in bacteria in a battery of genotoxicity tests conducted on microbial indicator organisms, in micronucleus tests in mice and hamsters, or in a sister chromatid exchange study in hamsters. No impairment of fertility was seen in male or female rats administered nicardipine at oral doses as high as 100 mg/kg/day (human equivalent dose about 16 mg/kg/day); thus the maximum recommended human oral dose).

13.2 Reproductive and Developmental Toxicology

Embryofetality, but no teratology, was seen at intrauterous doses of 10 mg nicardipine/kg/day in rats and 1 mg/kg/day in rabbits. These doses in the rat and rabbit are equivalent to human intrauterous doses of about 1.6 and 0.32 mg/kg/day, respectively. The total daily human dose delivered by a continuous intravenous infusion ranging from 1.2 to 6 mg/kg/day, depending on duration of different infusion rates ranging from 3 to 15 mg/kg as individual patients are titrated for optimal results, was also embryocidal when administered orally to pregnant Japanese white rabbits, during organogenesis, at 150 mg/kg/day (a dose associated with marked body weight gain suppression in the treated dose), but not at 50 mg/kg/day (human equivalent dose about 16 mg/kg/day or about 8 times the maximum recommended human oral dose). No adverse effects on the fetus were observed when New Zealand albino rabbits were treated orally, during organogenesis, with up to 100 mg/kg/day (a dose associated with significant mortality in the treated doe). In pregnant rats administered nicardipine orally at doses of up to 100 mg/kg/day (human equivalent dose about 16 mg/kg/day) there was no evidence of embryolethality or teratogenicity. However, dystopia, reduced birth weight, reduced neonatal survival and reduced neonatal weight gain were noted.

14 CLINICAL STUDIES

Effects in Hypertension

In patients with mild-to-moderate chronic stable essential hypertension, nicardipine hydrochloride injection (0.5 to 4 mg/hr) produced dose-dependent decreases in blood pressure, although only the decreases at 4 mg/hr were statistically different from placebo. At the end of a 48-hour infusion at 4 mg/hr, the decreases were 26 mmHg (17%) in systolic blood pressure and 21 mmHg (20%) in diastolic blood pressure. In other settings (e.g., patients with severe or postoperative hypertension), nicardipine hydrochloride injection (5 to 15 mg/hr) produced dose-dependent decreases in blood pressure. Higher infusion rates produced therapeutic responses more rapidly. The mean time to therapeutic response for severe hypertension, defined as diastolic blood pressure less than or equal to 95 mmHg or greater or equal to 25 mmHg decrease and systolic blood pressure less than or equal to 160 mmHg, was 77 ± 5 minutes. The average maintenance rate was 8.0 mg/hr. The mean time to therapeutic response for postoperative hypertension, defined as greater than or equal to 15% reduction in diastolic or systolic blood pressure, was 12 minutes. The average maintenance dose was 3 mg/hr.

16 HOW SUPPLIED/STORAGE AND HANDLING

16.1 How Supplied

Nicardipine hydrochloride in 0.9% sodium chloride injection is a color, clearless to yellow solution and is available in packages as follows:

NDC	Strength	Packaged
65145-197-01	20 mg in 200 mL (0.1 mg/mL)	1 single dose container
65145-198-01	40 mg in 200 mL (0.2 mg/mL)	1 single dose container

16.2 Storage and Handling

Store at 20° to 25°C (68° to 77°F) [see USP Controlled Room Temperature].

Nicardipine does not adversely affect the product, but exposure to elevated temperatures should be avoided. Protect from light.

Made in India.

CAPLIN STERILES

Distributed by:
Caplin Steriles USA Inc,
Hampton, NJ 08061
Code: TN Druugs/7N0000640S.

November 2025
22201534